

















- among adults from a developing country: the Sri Lanka diabetes and cardiovascular study. *Public Health Nutr* 16:1684-92.
- Kindermann R, Snell JL, 1980. Markov random fields and their applications. *Contemporary mathematics*. Vol 1. American Mathematical Society, Providence, RI, USA.
- Lang S, Brezger A, 2004. Bayesian P-splines. *J Comput Graph Stat* 13:183-212.
- Laraia BA, Blanchard SD, Karter AJ, Jones-Smith JC, Warton M, Kersten E, Jerrett M, Moffet HH, Adler N, Schillinger D, Kelly M, 2014. Spatial pattern of body mass index among adults in the diabetes study of Northern California (DISTANCE). *Int J Health Geogr* 13:48.
- Liese AD, Lawson A, Song H-R, Hibbert JD, Porter DE, Nichols M, Lamichhane AP, Dabelea D, Mayer-Davis EJ, Standiford D, Liu L, Hamman RF, D'Agostino RB Jr, 2010. Evaluating geographic variation in type 1 and type 2 diabetes mellitus incidence in youth in four US regions. *Health Place* 16:547-56.
- Liese AD, Puett RC, Lamichhane AP, Nichols MD, Dabelea D, Lawson AB, Porter DE, Hibbert JD, D'Agostino RB Jr, Mayer-Davis EJ, 2012. Neighborhood level risk factors for type 1 diabetes in youth: the SEARCH case-control study. *Int J Health Geogr* 11:1.
- Maty SC, Everson-Rose SA, Haan MN, Raghunathan TE, Kaplan GA, 2005. Education, income, occupation, and the 34-year incidence (1965-99) of Type 2 diabetes in the Alameda County Study. *Int J Epidemiol* 34:1274-81.
- Melidonis AM, Tournis SM, Kompoti MK, Lentzas IL, Roussou VR, Iraklianos SL, Michail IM, Mariolis AM, 2006. Increased prevalence of diabetes mellitus in a rural Greek population. *Rural Remote Health* 6:534.
- Mohan V, Mathur P, Deepa R, Deepa M, Shukla DK, Menon GR, Anand K, Desai NG, Joshi PP, Mahanta J, Thankappan KR, Shah B, 2008. Urban rural differences in prevalence of self-reported diabetes in India – the WHO-ICMR Indian NCD risk factor surveillance. *Diabetes Res Clin Pr* 80:159-68.
- Ning F, Pang ZC, Dong YH, Gao WG, Nan HR, Wang SJ, Zhang L, Ren J, Tuomilehto J, Hammar N, Malmberg K, Andersson SW, Qiao Q, Qingdao Diabetes Survey Group, 2009. Risk factors associated with the dramatic increase in the prevalence of diabetes in the adult Chinese population in Qingdao, China. *Diabetic Med* 26:855-63.
- Pearson JF, Bachireddy C, Shyamprasad S, Goldfine AB, Brownstein JS, 2010. Association between fine particulate matter and diabetes prevalence in the US. *Diabetes Care* 33:2196-201.
- Ramirez AG, Thompson IM, Vela L, 2008. *The South Texas health status review*. Springer, Amsterdam, The Netherlands.
- Schlundt DG, Hargreaves MK, McClellan L, 2006. Geographic clustering of obesity, diabetes, and hypertension in Nashville, Tennessee. *J Ambul Care Manage* 29:125-32.
- Sullivan PW, Morroto EH, Ghushchyan V, Wyatt HR, Hill JO, 2005. Obesity, inactivity, and the prevalence of diabetes and diabetes-related cardiovascular comorbidities in the US, 2000-2002. *Diabetes Care* 28:1599-603.
- US Census Bureau, 2014. American community survey information guide. Available from: <https://www.census.gov/programs-surveys/acs/about/information-guide.html>
- Voutilainen A, Hänninen S, Sherwood PR, Taipale H, Tolppanen AM, Vehviläinen-Julkunen K, 2015. Associations across spatial patterns of disease incidences, socio-demographics, and land use in Finland 1975-2010. *Scand J Public Health* 43:356-63.
- Zhou LL, AR, Adar SD, Park SK, 2014. Carbon dioxide emissions and change in prevalence of obesity and diabetes in the United States: an ecological study. *Environ Int* 73:111-6.

Non commercial use only